

Karow A, Reimer J, Schulz H, et al. Cost-utility analysis of 12 months Assertive Community Treatment as part of Integrated Care versus Standard Care in patients with schizophrenia treated with Quetiapine (ACCESS Trial). *J Clin Psychiatry* 2012; 73: 402-408.

Lambert M, Bock T, Schöttle D, et al. Assertive Community Treatment (ACT) as part of Integrated Care versus Standard Care: a 12-month trial in patients with first- and negatively selected multiple-episode schizophrenia-spectrum disorders treated with quetiapine IR. *J Clin Psychiatry* 2010; 71: 1313-23.

Lambert M, Bock T, Daubmann A, et al. The Hamburg-model of integrated care for patients with psychosis: Part 1. Rationale, treatment concept and results of the pre-study. *Psychiatr Prax* 2014a; 41: 257-65.

Lambert M, Schöttle D, Bock T, et al. Integrierte Versorgung nach dem Hamburger Modell für Patienten mit schweren psychotischen Erkrankungen: Behandlungsmodell, Patientencharakteristika, Therapieintensität und Effektivitätsergebnisse. *Psychotherapeut* 2014b; 59: 95–99.

Lambert M, Schöttle D, Ruppelt F, et al. Integrierte Versorgung für erst- und mehrfach erkrankte Patienten mit schweren psychotischen Erkrankungen: 3-Jahres Ergebnisse des Hamburger Modells. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2015a; 58: 408-19.

Lambert M, Schöttle D, Sengutta M, et al. Early Detection and Integrated Care in Adolescents and Young Adults with Severe Psychotic Illnesses. *Psychiatr Prax* 2015b;42 Suppl 1: 49-53.

Lambert M, Ruppelt F, Rohenkohl A, et al. Langzeittherapie in der Schizophrenie - Erfolgreiche psychosoziale Re-Integration durch Integrierte Versorgung. *Psychopharmakotherapie* 2016a; 23: 58–66.

Lambert M, Schöttle D, Sengutta M, et al. Early detection and integrated care for adolescents and young adults with severe psychotic disorders: rationale and design of the Integrated Care in Early Psychosis study (ACCESS III). *Early Interv Psychiatry*, 2016b Sep 12 [Epub ahead of print].

Lambert M, Schöttle D, Ruppelt F, et al. Early detection and integrated care for adolescents and young adults with psychotic disorders: the ACCESS III study. *Acta Psychiatr Scand*, 2017a Aug;136(2):188-200.

Schöttle D, Bock T, Meigel-Schleiff C, et al. Translating research into clinical practice: effectiveness of Integrated Care (IC) including therapeutic Assertive Community Treatment (ACT) in severe and persistent schizophrenia-spectrum and bipolar I disorders - a 24 months follow-up study (ACCESS II study). *J Clin Psychiatry* 2014; 75: 1371-1379.

Schöttle D, Ruppelt F, Karow A, et al. Hometreatment – Aufsuchende Behandlung im Hamburger Modell der Integrierten Versorgung. *Psychotherap Psych Med* 2015; 65: 140-145.